


SAFE MALE CIRCUMCISION (SMC)

Know your Facts











INTRODUCTION TO SAFE MALE CIRCUMCISION


- Hello and thank you for attending this session on SMC
- © I will ask a few questions throughout this session
- © Today we are going to discuss the benefits and implications of SMC
- © Feel free to ask questions at any time if you do not understand what I am saying
- © What do you know about safe male circumcision and HIV?

WHAT CAN YOU SEE?





ASK CLIENT WHAT HE CAN SEE



Coronal sulcus

Coronal rim

Urethral opening or Meatus

B


Circumcised penis

Uncircumcised Penis

Circumcised Penis




ASK CLIENTS WHAT THEY UNDERSTAND SAFE MALE CIRCUMCISION TO BE




- ☼ It was common in Botswana as part of the Bogwera ceremony until the colonial rulers stopped the practice in the early 20th Century
- SMC is the surgical removal of the foreskin covering the end of the penis so that the glans (or knob) is permanently exposed. SMC should be carried out in a sterile health facility by a trained doctor
- There are now specific health reasons why more and more men are choosing to do safe male circumcision





The goalkeeper needs the help of the team defenders to keep the ball out of the goal, just like a circumcised man needs the other defenders to prevent HIV infection

THE DEFENDERS



KNOW YOUR HIV STATUS

ALWAYS USE A CONDOM

STICK TO 1 PARTNER ABSTAIN FROM SEX


EXPLAIN THE DEFENDERS TO CLIENTS

The defenders of HIV are:

- © Testing for HIV regularly
- © Always using a condom correctly and consistently
- © Being faithful to one partner
- Abstaining from sex



DISCUSS THE BENEFITS OF SAFE MALE CIRCUMCISION



- © SMC increases penile hygiene a circumcised penis is easier to keep clean thereby reducing the risk of inflammation under the foreskin
- A circumcised man is also less likely to get cancer of the penis
- A circumcised man is less likely to get infected with the Human Papillomavirus (HPV), a virus that causes cervical cancer in women
- SMC reduces the risk of urinary tract infections (UTI's) in infants



EXPLAIN WHAT CLIENT SHOULD DO BEFORE THE PROCEDURE



- A qualified health professional or counsellor must be consulted before the procedure as they will explain about the SMC procedure and prepare you with all the facts necessary
- An HIV test should be carried out before the procedure because SMC is meant for HIV prevention in both partners
- © Circumcision as an HIV prevention strategy is not recommended for HIV-positive men as they are already infected. Circumcision cannot stop an HIV-positive man infecting or re-infecting a partner. Wound healing may take longer in HIV-positive men
- SMC has other health benefits for HIV-positive men and a session with a health care worker is recommended to understand more



DISCUSS THE RECOVERY PROCESS WITH CLIENTS

- © Complete recovery will take you about 6 weeks
- After the procedure you should return home and rest for one or two days as this will help the wound to heal
- You may bathe on the day after the surgery but must not let the dressing get wet
- © You should not have sexual intercourse or masturbate for 6 weeks. The wound must heal completely before sex



EXPLAIN WHAT THE CLIENT SHOULD EXPECT AFTER THE PROCEDURE

- © The risks to you, in getting circumcised by a qualified doctor in a sterile health facility, are very low
- However, you should expect a little pain or swelling where the wound is. Watch that it does not get worse, a small amount of pain that does not get worse is normal
- © Take the medication prescribed to you by your doctor
- © Look out for the following signs after the operation:
 - -Bleeding that does not stop or gets worse
 - -Severe pain
 - -Difficulty or inability to pass urine
 - -Pus discharging from the wound
 - -Increased swelling
 - -A fever within one week of surgery
 - -Severe lower abdominal pain
- If you experience any of these symptoms return to the health care facility immediately for further help



INFORM CLIENT WHEN HE CAN START HAVING SEX AGAIN

- © The doctor will advise you about your personal healing progress during your follow up visits
- Whilst you are recovering from the procedure do not have sexual intercourse or masturbate for 6 weeks. The wound must heal completely before resuming sexual activity. Having sex before the wound heals completely increases the risk of HIV infection
- Any injury to the penis can create an entry point for HIV
- The speed of recovery depends on resting, caring for the wound, abstaining from sex and returning to the clinic if any problems arise
- You may resume your normal sexual activity only after you receive confirmation of healing from your doctor




EXPLAIN WHAT IMPACT SMC HAS ON THE CLIENT'S SEX LIFE

- © The impact will vary from person to person as sexual pleasure depends on a large range of factors such as age, alcohol consumption, fatigue and the feelings sexual partners have towards each other
- © There is no scientific evidence that SMC either increases or decreases sexual pleasure for either partner




INFORM CLIENT HOW HE CAN PROTECT HIMSELF AND HIS PARTNER AFTER SMC

- Remember that a circumcised man can still get HIV because SMC does not provide complete protection.
- After circumcision you must continue to protect youself by;
 - -Testing for HIV regularly
 - -Always using a condom
 - -Being faithful to one partner
 - -Abstaining from sex



TELL CLIENTS WHAT TO SAY TO OTHERS WHO WANT TO GET CIRCUMCISED

- © You can explain that SMC can decrease their risk of HIV infection but it is only **partially protective**
- You can encourage them to go to a trained and experienced health care provider who offers counselling and HIV testing so that they can learn the benefits of this procedure
- You can also explain how a circumcised man is less likely to get infected by the Human Papillomavirus (HPV), a virus that causes cervical cancer in women
- © Tell others that getting circumcised shows responsibility because it proves that they are taking a measure against HIV



REVIEW WHAT WAS LEARNT TODAY

- SMC does not provide complete protection against HIV, so circumcised men CAN still get HIV if they don't protect themselves
- © It lowers the risk of getting sexually transmitted infections (STI's)
- Circumcised men find it easier to maintain penile hygiene and a circumcised man is less likely to get infected by the Human Papillomavirus (HPV), a virus that cauese cervical cancer in women.

After circumcision, you must continue to protect yourself by

- -Testing for HIV regularly
- -Always using a condom
- -Being faithful to one partner
- -Abstaining from sex





- © Do you know your facts?
- © Do you have any concerns about the procedure?
- © So do you want to get circumcised now that you know the facts?

FOR MORE INFORMATION
PLEASE CONTACT THE DEPARTMENT OF
HIV/AIDS PREVENTION AND CARE ON
0800 600 700 / 3632237 / 711600







SAFE MALE CIRCUMCISION (SMC)

Know your Facts