

Table 2. Numbers of male circumcisions performed, achievement and potential infections averted in eastern and southern Africa, 2010

Countries	Number of MCs done by MMC programme by calendar year				Estimated number of MCs needed to reach 80% prevalence*	Potential infections averted by scaling up MC to reach 80% prevalence* in 5 years		Achievement towards estimated number of MCs needed (%)
	2008	2009	2010	Total		Number	%	
Botswana	0	5424	5773	11 197	345 244	62 773	28	3.2
Ethiopia	0	769	2689	3458	40 000	1479	17	8.6
Kenya	11 663	80 719	139 905 <sup>a</sup>	232 287 <sup>a</sup>	860 000 <sup>a*</sup>	73 420 <sup>**</sup>	16 <sup>**</sup>	27.0*
Lesotho	0	0	Unknown <sup>b</sup>	Unknown <sup>b</sup>	376 795	106 427	34	0.06
Malawi	589 <sup>c</sup>	1234 <sup>c</sup>	1296 <sup>c</sup>	3119 <sup>c</sup>	2 101 566	240 685	28	0.1
Mozambique	0	100	7633	7733	1 059 104	215 861	13	0.7
Namibia	0	224	1763	1987	330 218	18 373	25	0.6
Rwanda	0	0	1694	1694	1 746 052	56 840	29	0.1
South Africa	5 190 <sup>d</sup>	9168 <sup>d</sup>	131 117 <sup>d</sup>	145 475 <sup>d</sup>	4 333 134	1 083 869	20	3.4
Swaziland	1 110	4336	18 869	24 315	183 450	56 810	34	13.3
Tanzania	0	1033	18 026 <sup>e</sup>	19 059 <sup>e</sup>	1 373 271	202 900	9	1.4
Uganda	0	0	9052	9052	4 245 184	339 524	25	0.2
Zambia	2758	17 180	61 911	81 849	1 949 292	339 632	30	4.2
Zimbabwe	0	2801	11 176	13 977	1 912 595	565 751	42	0.7
<b>Total</b>	<b>21 310</b>	<b>122 988</b>	<b>410 904</b>	<b>555 202</b>	<b>20 855 905*</b>	<b>3 364 345</b>	<b>22</b>	<b>2.7</b>

\* Estimated number of MCs needed is based on a target of 80% prevalence among males 15 - 49 years old; except in Kenya where it is based on the national goal of 94% of males 15 - 49 years old.

\*\* Nyanza Province only

Source: PEPFAR Male Circumcision Technical Working Group unless otherwise indicated, <sup>a</sup> Kenya NASCOP, <sup>b</sup> Lesotho MOH, <sup>c</sup> Malawi MOH, <sup>d</sup> South Africa NDOH, <sup>e</sup> Tanzania MOH