

Zimbabwe

2020

Based on most recent available data.

Version

2.01.

Output (coverage)

Outcome (service use/behaviour)

Impact

Summary

Output (coverage)		Outcome (service use/behaviour)		Impact		Summary														
Condoms																				
Number of condoms distributed and sold / year (in millions)	95	Condom use with non-regular partners (%)	Women 15-49	65	New HIV infections (ages 15+, in 1000s)	<table border="1"> <tr><td>AGYW&MPs</td><td>4</td></tr> <tr><td>SW</td><td>6</td></tr> <tr><td>MSM</td><td>5</td></tr> <tr><td>PWID</td><td>id</td></tr> <tr><td>Condoms</td><td>8</td></tr> <tr><td>VMMC</td><td>6</td></tr> <tr><td>PrEP</td><td>5</td></tr> </table>	AGYW&MPs	4	SW	6	MSM	5	PWID	id	Condoms	8	VMMC	6	PrEP	5
AGYW&MPs	4																			
SW	6																			
MSM	5																			
PWID	id																			
Condoms	8																			
VMMC	6																			
PrEP	5																			
Number of condoms distributed and sold per man 15-64 /year	20	Men 15-49	82																	
% of condom distribution need met for men 15-64	83																			
Voluntary medical male circumcision (VMMC)																				
Number of VMMCs performed / year (in thousands)	355	% of 2020 VMMC target achieved	Men	42																
% of annual VMMC target achieved	68	National male circumcision prevalence (%)	Men 15-24	19																
			Men 15-49	14																
ARV-based prevention																				
Composite PrEP score (0-10)	5	% of national PrEP need met	All pop.	id	<table border="1"> <tr><td>HIV prevalence</td><td>6</td></tr> </table>	HIV prevalence	6													
HIV prevalence	6																			
Number of people who received PrEP at least once in the past 12 months	8351	% of PLHIV virally suppressed	Women 15+	79																
% of PLHIV on ART	85	% of PLHIV virally suppressed	Men 15+	69																
Key populations																				
<i>Sex workers (SW)</i>																				
Population size estimate for sex workers (in thousands)	45	Condom use / last paid sex (%)	Sex workers	75	Sex workers <25 years	25														
% of SWs who received at least two HIV prevention interventions (past 3 mo)	44	Condom use / last paid sex (%)	Men 15-49	90	Sex workers all ages	42														
Prevention strategy includes core elements of SW prevention package	All	% of PLHIV on ART	SWs LHIV	78																
<i>Gay men and other men who have sex with men (MSM)</i>																				
Population size estimate for men who have sex with men (in thousands)	16	Condom use / last anal sex (%)	MSM	69	MSM <25 years	9														
% of MSM who received at least two HIV prevention interventions (past 3 mo)	26	% of PLHIV on ART	MSM LHIV	94	MSM all ages	21														
Prevention strategy includes core elements of MSM prevention package	< Half																			
<i>People who inject drugs (PWID)</i>																				
Population size estimate for people who inject drugs (in thousands)	id	% with safe injecting practices	PWID	id	PWID <25 years	id														
% of PWID who received at least two HIV prevention interventions (past 3 mo)	id	% on opioid substitution therapy	PWID	id	PWID all ages	id														
Prevention strategy includes core elements of PWID harm reduction package	None	% of PLHIV on ART	PWID LHIV	id																
<i>Structural barriers and enablers</i>																				
Criminalization of selling sex	Yes	% of people who avoided health care because of stigma and discrimination	Sex workers	39																
Criminalization of same sex relations	Yes		MSM	8																
Criminalization of drug use/consumption or possession for personal use	Yes		PWID	id																
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings																				
% of priority locations/districts with dedicated programs for AGYW & partners	17	Condom use with non-regular partners (%)	YW 15-24	54	Young women 15-24	5.9														
% of girls who completed lower secondary education	74		YM 15-24	81	Young men 15-24	3.3														
Educational policies on HIV & sexuality education (secondary school)	Yes	% who had multiple sexual partners	Men 15-49	14	Adults 15-49	12.8														
Laws requiring parental consent for adolescents to access SRH services	Yes	% of women experienced physical or sexual violence from husband/partner	YW 15-19	31																
Provider-initiated condom promotion integrated into SRH services	Yes		Women 15-49	19																
HIV testing services integrated with SRH services	Yes																			

Acronyms na ... not applicable;

PLHIV...people living with HIV; LHIV...living with HIV;YW...young women;YM...young men;

OST ... opioid substitution therapy; SRH ... sexual and reproductive health.

Very good

Good

Medium

Low

Very low

id Insufficient data