Lesotho	2020	Based on most recent available data.	Version	2.01.		
Output (coverage)		Outcome (service use/beha	aviour)	$\rightarrow$	P Impact	Summary
Condoms					New HIV infections (ages 15+, in 1000	Scores
Number of condoms distributed and sold / year (in millions)	38	Condom use with non-regular partners (%)	Women 15-49	66	17	AGYW&MPs
Number of condoms distributed and sold per man 15-64 /year	56	condom use with non regular partners (70)	Men 15-49	73	18	8
% of condom distribution need met for men 15-64	100				16 14	SW
Voluntary medical male circumcision (VMMC)					12 10	4
Number of VMMCs performed / year (in thousands)	34	% of 2020 VMMC target achieved	Men	58	10	MSM
% of annual VMMC target achieved	88	National male circumcision prevalence (%)	Men 15-24	70	8	5
			Men 15-49	69	6	4 PWID
ARV-based prevention					4	♦ id
Composite PrEP score (0-10)	10	% of national PrEP need met	All pop.	id	2	Condoms
Number of people who received PrEP at least once in the past 12 months	35478	% of PLHIV virally surpressed	Women 15+	66	2010 2015	2020
% of PLHIV on ART	65	% of PLHIV virally surpressed	Men 15+	54		VMMC
Key populations					HIV preval	ence 8
Sex workers (SW)						PrEP
Population size estimate for sex workers (in thousands)	8	Condom use / last paid sex (%)	Sex workers	62	Sex workers <25 years id	10
% of SWs who received at least two HIV prevention interventions (past 3 mo)	27	Condom use / last paid sex (%)	Men 15-49	90	Sex workers all ages 72	
Prevention strategy includes core elements of SW prevention package	> Half	% of PLHIV on ART	SWs LHIV	id		
Gay men and other men who have sex with men (MSM)						
Population size estimate for men who have sex with men (in thousands)	6	Condom use / last anal sex (%)	MSM	46	MSM <25 years id	
% of MSM who received at least two HIV prevention interventions (past 3 mo)	49	% of PLHIV on ART	MSM LHIV	id	MSM all ages 33	
Prevention strategy includes core elements of MSM prevention package	> Half					_
People who inject drugs (PWID)						
Population size estimate for people who inject drugs (in thousands)	id	% with safe injecting practices	PWID	id	PWID <25 years id	
% of PWID who received at least two HIV prevention interventions (past 3 mo)	id	% on opioid substitution therapy	PWID	id	PWID all ages id	
Prevention strategy includes core elements of PWID harm reduction package	None	% of PLHIV on ART	PWID LHIV	id		
Structural barriers and enablers						
Criminalization of selling sex	Yes		Sex workers	8		
Criminalization of same sex relations	No	% of people who avoided health care because of stigma and discrimination	MSM	8		
Criminalization of drug use/consumption or possession for personal use	Yes		PWID	id		
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings						
% of priority locations/districts with dedicated programs for AGYW & partners	100	Condom use with non-regular partners (%)	YW 15-24	82	Young women 15-24 10.	L
% of girls who completed lower secondary education	46		YM 15-24	79	Young men 15-24 4.5	
Educational policies on HIV & sexuality education (secondary school)	Yes	% who had multiple sexual partners	Men 15-49	27	Adults 15-49 22.	3
Laws requiring parental consent for adolescents to access SRH services	Yes	% of women experienced physical or sexual	YW 15-19	id	1	
Provider-initiated condom promotion integrated into SRH services	No/id	violence from husband/partner	Women 15-49	1		
HIV testing services integrated with SRH services	Yes					
Acronyms na not applicable;		Very good			Low	
PLHIVpeople living with HIV; LHIVliving with HIV;YWyoung women;YMyoung men;		Good			Very low	
OST opioid substitution therapy; SRH sexual and reproductive health.		Medium		id	Insufficient data	