

# Mozambique

2020

Based on most recent available data.

Version

2.01.

## Output (coverage)

## Outcome (service use/behaviour)

## Impact

## Summary

Output (coverage)		Outcome (service use/behaviour)		Impact		Summary
<b>Condoms</b>						
Number of condoms distributed and sold / year (in millions)	96	Condom use with non-regular partners (%)	Women 15-49	42		<b>AGYW&amp;MPs</b> 4 <b>SW</b> 4 <b>MSM</b> id <b>PWID</b> 2 <b>Condoms</b> 4 <b>VMMC</b> 6 <b>PrEP</b> 3
Number of condoms distributed and sold per man 15-64 / year	15		Men 15-49	47		
% of condom distribution need met for men 15-64	44					
<b>Voluntary medical male circumcision (VMMC)</b>						
Number of VMMCs performed / year (in thousands)	391	% of 2020 VMMC target achieved	Men	41		
% of annual VMMC target achieved	67	National male circumcision prevalence (%)	Men 15-24	66		
			Men 15-49	63		
<b>ARV-based prevention</b>						
Composite PrEP score (0-10)	3	% of national PrEP need met	All pop.	id		
Number of people who received PrEP at least once in the past 12 months	7434	% of PLHIV virally suppressed	Women 15+	53		
% of PLHIV on ART	60	% of PLHIV virally suppressed	Men 15+	35		
<b>Key populations</b>						
<i>Sex workers (SW)</i>						
Population size estimate for sex workers (in thousands)	93	Condom use / last paid sex (%)	Sex workers	id	Sex workers <25 years	id
% of SWs who received at least two HIV prevention interventions (past 3 mo)	51	Condom use / last paid sex (%)	Men 15-49	31	Sex workers all ages	id
Prevention strategy includes core elements of SW prevention package	> Half	% of PLHIV on ART	SWs LHIV	id		
<i>Gay men and other men who have sex with men (MSM)</i>						
Population size estimate for men who have sex with men (in thousands)	41	Condom use / last anal sex (%)	MSM	id	MSM <25 years	id
% of MSM who received at least two HIV prevention interventions (past 3 mo)	4	% of PLHIV on ART	MSM LHIV	id	MSM all ages	id
Prevention strategy includes core elements of MSM prevention package	> Half					
<i>People who inject drugs (PWID)</i>						
Population size estimate for people who inject drugs (in thousands)	14	% with safe injecting practices	PWID	id	PWID <25 years	id
% of PWID who received at least two HIV prevention interventions (past 3 mo)	15	% on opioid substitution therapy	PWID	id	PWID all ages	id
Prevention strategy includes core elements of PWID harm reduction package	Some	% of PLHIV on ART	PWID LHIV	id		
<i>Structural barriers and enablers</i>						
Criminalization of selling sex	No	% of people who avoided health care because of stigma and discrimination	Sex workers	id		
Criminalization of same sex relations	No		MSM	id		
Criminalization of drug use/consumption or possession for personal use	Yes		PWID	id		
<b>Adolescent girls, young women (AGYW) &amp; partners in high-HIV incidence settings</b>						
% of priority locations/districts with dedicated programs for AGYW & partners	42	Condom use with non-regular partners (%)	YW 15-24	51	Young women 15-24	7.1
% of girls who completed lower secondary education	10		YM 15-24	48	Young men 15-24	2.8
Educational policies on HIV & sexuality education (secondary school)	Yes	% who had multiple sexual partners	Men 15-49	21	Adults 15-49	12.4
Laws requiring parental consent for adolescents to access SRH services	No	% of women experienced physical or sexual violence from husband/partner	YW 15-19	10		
Provider-initiated condom promotion integrated into SRH services	Yes		Women 15-49	16		
HIV testing services integrated with SRH services	Partial					

**Acronyms** na ... not applicable;

PLHIV...people living with HIV; LHIV...living with HIV; YW...young women; YM...young men;

OST ... opioid substitution therapy; SRH ... sexual and reproductive health.

Very good

Good

Medium

Low

Very low

id Insufficient data