

Output (coverage)

Outcome (service use/behaviour)

Impact

Summary

Output (coverage)		Outcome (service use/behaviour)		Impact		Summary
Condoms						
Number of condoms distributed and sold / year (in millions)	29	Condom use with non-regular partners (%)	Women 15-49	59		Scores AGYW&MPs 5 SW 6 MSM id PWID id Condoms 8 VMMC 5 PrEP 8
Number of condoms distributed and sold per man 15-64 /year	37		Men 15-49	70		
% of condom distribution need met for men 15-64	100					
Voluntary medical male circumcision (VMMC)						
Number of VMMCs performed / year (in thousands)	41	% of 2020 VMMC target achieved	Men	34	HIV prevalence Sex workers <25 years 14 Sex workers all ages 28 MSM <25 years 4 MSM all ages 9 PWID <25 years id PWID all ages id	
% of annual VMMC target achieved	55	National male circumcision prevalence (%)	Men 15-24	27		
			Men 15-49	24		
ARV-based prevention						
Composite PrEP score (0-10)	8	% of national PrEP need met	All pop.	id		
Number of people who received PrEP at least once in the past 12 months	10583	% of PLHIV virally suppressed	Women 15+	83		
% of PLHIV on ART	85	% of PLHIV virally suppressed	Men 15+	69		
Key populations						
<i>Sex workers (SW)</i>						
Population size estimate for sex workers (in thousands)	8	Condom use / last paid sex (%)	Sex workers	86		
% of SWs who received at least two HIV prevention interventions (past 3 mo)	60	Condom use / last paid sex (%)	Men 15-49	67		
Prevention strategy includes core elements of SW prevention package	> Half	% of PLHIV on ART	SWs LHIV	34		
<i>Gay men and other men who have sex with men (MSM)</i>						
Population size estimate for men who have sex with men (in thousands)	7	Condom use / last anal sex (%)	MSM	id		
% of MSM who received at least two HIV prevention interventions (past 3 mo)	64	% of PLHIV on ART	MSM LHIV	15		
Prevention strategy includes core elements of MSM prevention package	> Half					
<i>People who inject drugs (PWID)</i>						
Population size estimate for people who inject drugs (in thousands)	id	% with safe injecting practices	PWID	id		
% of PWID who received at least two HIV prevention interventions (past 3 mo)	id	% on opioid substitution therapy	PWID	id		
Prevention strategy includes core elements of PWID harm reduction package	None	% of PLHIV on ART	PWID LHIV	id		
<i>Structural barriers and enablers</i>						
Criminalization of selling sex	Yes	% of people who avoided health care because of stigma and discrimination	Sex workers	id		
Criminalization of same sex relations	Yes		MSM	id		
Criminalization of drug use/consumption or possession for personal use	Yes		PWID	id		
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings						
% of priority locations/districts with dedicated programs for AGYW & partners	29	Condom use with non-regular partners (%)	YW 15-24	64	Young women 15-24	4.8
% of girls who completed lower secondary education	62		YM 15-24	78	Young men 15-24	2.7
Educational policies on HIV & sexuality education (secondary school)	Yes	% who had multiple sexual partners	Men 15-49	10	Adults 15-49	11.5
Laws requiring parental consent for adolescents to access SRH services	No	% of women experienced physical or sexual violence from husband/partner	YW 15-19	0		
Provider-initiated condom promotion integrated into SRH services	No/id		Women 15-49	1		
HIV testing services integrated with SRH services	Yes					

Acronyms na ... not applicable;

PLHIV...people living with HIV; LHIV...living with HIV; YW...young women; YM...young men;

OST ... opioid substitution therapy; SRH ... sexual and reproductive health.

Very good

Good

Medium

Low

Very low

id Insufficient data