Uganda	2020	Based on most recent available data.	Version	2.01.			
Output (coverage)		Outcome (service use/beha	aviour)		Impact		Summary
Condoms Number of condoms distributed and sold / year (in millions) Number of condoms distributed and sold per man 15-64 /year % of condom distribution need met for men 15-64	138 14 62	Condom use with non-regular partners (%)	Women 15-49 Men 15-49	38 62	New HIV infections (ages 1 80 72 70	5+, in 1000s)	Scores AGYW&MPs 4 SW
Voluntary medical male circumcision (VMMC)					60	48	5
Number of VMMCs performed / year (in thousands)	769	% of 2020 VMMC target achieved	Men	54	50 40	40	MSM
% of annual VMMC target achieved	84	National male circumcision prevalence (%)	Men 15-24 Men 15-49	49 43	30	18	3 PWID
ARV-based prevention					20	•	1
Composite PrEP score (0-10)	6	% of national PrEP need met	All pop.	id	10		Condoms
Number of people who received PrEP at least once in the past 12 months	38064	% of PLHIV virally surpressed	Women 15+	83			6
% of PLHIV on ART	84	% of PLHIV virally surpressed	Men 15+	68	2010 2015	2020	VMMC
Key populations						HIV prevalence	8
Sex workers (SW)							PrEP
Population size estimate for sex workers (in thousands)	174	Condom use / last paid sex (%)	Sex workers	69	Sex workers <25 yea	rs id	6
% of SWs who received at least two HIV prevention interventions (past 3 mo)	42	Condom use / last paid sex (%)	Men 15-49	73	Sex workers all ag	es 31	
Prevention strategy includes core elements of SW prevention package	All	% of PLHIV on ART	SWs LHIV	65			
Gay men and other men who have sex with men (MSM)							
Population size estimate for men who have sex with men (in thousands)	44	Condom use / last anal sex (%)	MSM	39	MSM <25 yea	rs id	
% of MSM who received at least two HIV prevention interventions (past 3 mo)	12	% of PLHIV on ART	MSM LHIV	66	MSM all ag	es 13	
Prevention strategy includes core elements of MSM prevention package	> Half						
People who inject drugs (PWID)							
Population size estimate for people who inject drugs (in thousands)	7	% with safe injecting practices	PWID	id	PWID <25 yea	rs 17	
% of PWID who received at least two HIV prevention interventions (past 3 mo)	8	% on opioid substitution therapy	PWID	id	PWID all ag	es 17	
Prevention strategy includes core elements of PWID harm reduction package	None	% of PLHIV on ART	PWID LHIV	78			
Structural barriers and enablers							
Criminalization of selling sex	Yes		Sex workers	id			
Criminalization of same sex relations	Yes	% of people who avoided health care because of stigma and discrimination	MSM	id			
Criminalization of drug use/consumption or possession for personal use	Yes		PWID	64			
Adolescent girls, young women (AGYW) & partners in high-HIV incidence settings							
% of priority locations/districts with dedicated programs for AGYW & partners	31	Condom use with non-regular partners (%)	YW 15-24	44	Young women 15-2	24 2.8	
% of girls who completed lower secondary education	36		YM 15-24	63	Young men 15-2	24 1.2	
Educational policies on HIV & sexuality education (secondary school)	Yes	% who had multiple sexual partners	Men 15-49	21	Adults 15-4	19 5.8	
Laws requiring parental consent for adolescents to access SRH services	No	% of women experienced physical or sexual	YW 15-19	31			
Provider-initiated condom promotion integrated into SRH services	Yes	violence from husband/partner	Women 15-49	50			
HIV testing services integrated with SRH services	Yes		I				
Acronyms na not applicable;		Very good			Low		
PLHIVpeople living with HIV; LHIVliving with HIV;YWyoung women;YMyoung men;		Good			Very low		
OST opioid substitution therapy; SRH sexual and reproductive health.		Medium		id	Insufficient data		