

# Zambia

2020

Based on most recent available data.

Version

2.01.

## Output (coverage)

## Outcome (service use/behaviour)

## Impact

## Summary

Output (coverage)		Outcome (service use/behaviour)		Impact	Summary	
<b>Condoms</b>						
Number of condoms distributed and sold / year (in millions)	62	Condom use with non-regular partners (%)	Women 15-49	35	<p>New HIV infections (ages 15+, in 1000s)</p>	
Number of condoms distributed and sold per man 15-64 / year	13		Men 15-49	62		
% of condom distribution need met for men 15-64	51					
<b>Voluntary medical male circumcision (VMMC)</b>						
Number of VMMCs performed / year (in thousands)	550	% of 2020 VMMC target achieved	Men	71		
% of annual VMMC target achieved	100	National male circumcision prevalence (%)	Men 15-24	37		
			Men 15-49	65		
<b>ARV-based prevention</b>						
Composite PrEP score (0-10)	6	% of national PrEP need met	All pop.	id		
Number of people who received PrEP at least once in the past 12 months	23250	% of PLHIV virally suppressed	Women 15+	82		
% of PLHIV on ART	85	% of PLHIV virally suppressed	Men 15+	71		
<b>Key populations</b>						
<i>Sex workers (SW)</i>						
Population size estimate for sex workers (in thousands)	134	Condom use / last paid sex (%)	Sex workers	79	Sex workers <25 years id	
% of SWs who received at least two HIV prevention interventions (past 3 mo)	47	Condom use / last paid sex (%)	Men 15-49	56	Sex workers all ages 49	
Prevention strategy includes core elements of SW prevention package	> Half	% of PLHIV on ART	SWs LHIV	86		
<i>Gay men and other men who have sex with men (MSM)</i>						
Population size estimate for men who have sex with men (in thousands)	68	Condom use / last anal sex (%)	MSM	id	MSM <25 years id	
% of MSM who received at least two HIV prevention interventions (past 3 mo)	13	% of PLHIV on ART	MSM LHIV	id	MSM all ages id	
Prevention strategy includes core elements of MSM prevention package	> Half					
<i>People who inject drugs (PWID)</i>						
Population size estimate for people who inject drugs (in thousands)	27	% with safe injecting practices	PWID	id	PWID <25 years id	
% of PWID who received at least two HIV prevention interventions (past 3 mo)	id	% on opioid substitution therapy	PWID	id	PWID all ages id	
Prevention strategy includes core elements of PWID harm reduction package	None	% of PLHIV on ART	PWID LHIV	id		
<i>Structural barriers and enablers</i>						
Criminalization of selling sex	Partial	% of people who avoided health care because of stigma and discrimination	Sex workers	id		
Criminalization of same sex relations	Yes		MSM	id		
Criminalization of drug use/consumption or possession for personal use	Yes		PWID	id		
<b>Adolescent girls, young women (AGYW) &amp; partners in high-HIV incidence settings</b>						
% of priority locations/districts with dedicated programs for AGYW & partners	16	Condom use with non-regular partners (%)	YW 15-24	34	Young women 15-24 5.5	
% of girls who completed lower secondary education	48		YM 15-24	49	Young men 15-24 2.6	
Educational policies on HIV & sexuality education (secondary school)	Yes	% who had multiple sexual partners	Men 15-49	15	Adults 15-49 11.5	
Laws requiring parental consent for adolescents to access SRH services	Yes	% of women experienced physical or sexual violence from husband/partner	YW 15-19	27		
Provider-initiated condom promotion integrated into SRH services	Yes		Women 15-49	25		
HIV testing services integrated with SRH services	Partial					

**Acronyms** na ... not applicable;

PLHIV...people living with HIV; LHIV...living with HIV; YW...young women; YM...young men;

OST ... opioid substitution therapy; SRH ... sexual and reproductive health.

Very good

Good

Medium

Low

Very low

id Insufficient data

### Scores

AGYW&MPs

3

SW

5

MSM

id

PWID

id

Condoms

5

VMMC

9

PrEP

6

HIV prevalence